Why should I care about trying to prevent an ileus?

- An ileus can cause you pain or discomfort
- You may have nausea or vomiting. An ileus can increase your risk for lung problems
- Some patients may have to go to the Emergency Department or be readmitted to the hospital for their ileus

That sounds serious!

What can I do to prevent this?

To watch a video on education and expectations post-surgery, scan the QR code (below) with the camera on your smartphone device or visit www.musicurology.com/rpvideo/



- 1. Open the camera app on your smartphone device.
- 2. Scan the QR code (left).
- 3. Click on the YouTube banner link to begin the video.



For additional information regarding this informational pamphlet, please contact us at:

MUSIC Coordinating Center

2800 Plymouth Rd. Building 16 Ann Arbor, MI 48109

(855) 456 - 2035 www.musicurology.com

Copyright 2018.
For questions about the Michigan Urological Surgery Improvement Collaborative, please contact us at musicurology@umich.edu

MUSIC is sponsored by:



Blue Cross Blue Shield of Michigan

A nonprofit corporation and independent licensee of the Blue Cross and Blue Shield Association

You and your doctor decided on surgery to treat your prostate cancer.

- As with any surgery, there are potential complications
- You and your doctor may have already discussed erectile dysfunction and incontinence
- One problem that can make you return to the hospital is an ileus



Preventing ILEUS:


A common reason for an unplanned return to the hospital


What is ileus?


- Ileus is severe constipation
- It is when you cannot pass gas or have a bowel movement
- Ileus is a common problem after prostate surgery

Why may ileus be related to my surgery?

- The intestines can go to sleep after abdominal or pelvic surgery
- Narcotic (opioid) pain medication can make ileus worse







Before surgery



Do not eat or drink anything unless directed by your doctor or nurse.

After surgery



Walk around and try to pass gas.

The day **AFTER** surgery:



Try to get up every 2 hours to walk around. Walking around may help you pass gas.

When home with your catheter



Continue to walk around, care for your catheter, and follow a small portion soft diet.

When you're home, for the next 30 days:



- Gradually build your diet from small to normal portions.
- Follow a soft diet until regular bowel movements return.
- Stay active. Continue walking, but avoid strenuous, weight bearing exercise.





• If sitting for longer than 1 hour, take a short walk at least once per hour while you are awake.









What will it look like if I have an ileus?

- If you develop an ileus, you will most likely be very bloated/constipated and have a lot of abdominal and pelvic pain/discomfort.
- You may not be able to pass gas or have difficulty passing a bowel movement.



Get Help!

When should I call someone?

Severe pain, fever, chills, nausea, vomiting, inability to keep down food and water

Who should I call?

My doctor's office

What should I do if I cannot reach anyone?

Seek care in an emergency department or urgent care

For more information, please visit: www.musicUrology.com/NOTES